

URINE COLLECTION FOR BACTERIAL CULTURE

DEAR PATIENT,


Your doctor has ordered a laboratory test that requires the collection of a midstream urine sample.

It is important to collect the sample correctly.

This guide explains how to collect the urine sample, label it, and store it properly until delivery to the laboratory.

From the laboratory you will need:

- Single-use sterile urine collection container
- 10 mL sterile syringe
- Green-cap urine container with stabilizer



HOW IS A MIDSTREAM URINE SAMPLE COLLECTED?

What should you pay attention to:

- If possible, the midstream urine sample should be collected from the first urination after waking up in the morning. Otherwise, there should be at least a 4-hour interval between urinations
- Sexual intercourse should be avoided for 24 hours before sample collection.
- First, label the urine container on the side with your name, date of birth, and date of collection, and place it nearby with the lid within easy reach.
- Wash your hands thoroughly with soap and water, then dry them with a clean towel.




WHAT HAPPENS AFTER URINE COLLECTION?

- Hold the green-capped container upright and remove the cap. Lock the plunger in its final position. Using a sterile syringe, transfer 10 mL of urine from the single-use sterile urine collection container into the open part of the green-capped container. Close the cap tightly and shake the container gently.
- Bring the green-capped container to the laboratory within a maximum of one hour.

COLLECTION INSTRUCTIONS:

1. Wash your genital area

Women: With one hand, spread the labia apart and wash the area thoroughly using water only (without soap). Then dry the area with a clean towel. Keep the labia separated until urine collection is completed. Urine collection during menstruation is not recommended. If necessary, a tampon should be used.



Men: Retract the foreskin completely and clean the glans thoroughly using water only (without soap). Then dry the area carefully with a clean towel. Keep the foreskin retracted until urine collection is completed.

2. Collect the Midstream Urine Sample

Let the urine to flow into the toilet for approximately 3 seconds. Then collect approximately 20–30 mL of midstream urine in the urine container without interrupting urination. The collection container should not come into contact with the body. Allow the remaining urine to flow into the toilet

